

Rational pencils of cubics with 6 real base points

S  verine Fiedler-Le Touz  

October 29, 2012

Abstract

Let six points $1, \dots, 6$ lie in general position in the real projective plane and consider the pencil of nodal cubics based at these points, with node at one of them, say 1. This pencil has five reducible cubics. We call combinatorial cubic a topological type (cubic, base points) and combinatorial pencil the cyclic sequence of five combinatorial reducible cubics. Up to the action of the symmetric group $\mathcal{S}_5$ on $\{2, \dots, 6\}$, there are seven possible combinatorial pencils with node at 1. Consider now the set of six pencils obtained, making the node to be $1, \dots, 6$. Up to the action of $\mathcal{S}_6$ on $\{1, \dots, 6\}$, there are four lists of six combinatorial pencils.

Let $1, \dots, 6$ be six points in the real projective plane and consider the pencil of nodal cubics determined by these points with node at one of them, say 1. This pencil has five reducible cubics plus, possibly, some cuspidal cubics. We ignore the latter and consider the sequence of reducible cubics. Let us call combinatorial cubic a topological type (cubic, base points) and combinatorial pencil the cyclic sequence of five combinatorial reducible cubics, they are of the form $1m \cup 1ijkl$, where $\{i, j, k, l, m\} = \{2, 3, 4, 5, 6\}$. We shall prove the following:

Theorem 1 *Let $1, \dots, 6$ be six points in $\mathbb{RP}^2$. Up to the action of $\mathcal{S}_5$ on $\{2, \dots, 6\}$, there are seven possible combinatorial pencils of nodal cubics based at these six points with node at 1.*

Up to the action of $\mathcal{S}_6$ on $\{1, \dots, 6\}$, there are four possible lists of six combinatorial pencils of nodal cubics based at these six points with respective nodes at $1, \dots, 6$.

Proof: The conic and the 10 lines determined by five of the points divide the plane in 36 zones. Up to the action of $\mathcal{S}_5$ on $\{2, \dots, 6\}$, we may assume that the six points are disposed as shown in Figure 1, where 2, 5, 3, 6, 4 lie in this ordering on a conic, and 1 is in one of the zones $A, \dots, G$. Let $1 \in A$. We perform a cremona transformation $cr : (x_0; x_1; x_2) \rightarrow (x_1x_2; x_0x_2; x_0x_1)$ with base points 1, 4, 5. Let us denote the respective images of the lines 14, 15, 45 by 5, 4, 1. For the other points, we keep the same notation as before cr . The three conics 14365, 31452, 61542 are mapped onto three lines 36, 23 and 26, see upper part of Figure 2. After cr , let us consider the pencil of conics 1236, and five particular conics of this pencil: the three double lines and the two conics passing respectively through 4 and 5. The cyclic ordering of these conics in the pencil is easily determined. Perform the cremona transformation back. The pencil of conics 1236 is mapped onto the pencil of nodal cubics based at 1, $\dots, 6$ with node at 1. The five particular conics are mapped onto the five reducible cubics. We repeat this procedure for the various positions of $1 \in B, \dots, G$, with a cremona transformation based either in 145 or in 136, see Figures 2-5. The sequences of five particular conics are displayed in Figure 6. The sequences of five reducible cubics for each case $A, \dots, G$ are shown in Figure 7, the pencils are drawn in Figures 8-11. More precisely, we have represented the successive types of combinatorial cubics in the five portions bounded by the reducible cubics. The upper middle pictures of each pencil B, C, G are not quite correct: in the actual pencils, the portion starts and finishes with cubics having a loop containing no base point other than 1. Inbetween, there is a pair of cuspidal cubics, and in the middle, cubics with an isolated node. We have represented only the latter.

Consider any one of these seven pencils. We observe the following properties: the cyclic ordering of the points 2, $\dots, 6$ is the same on each of the five types of non-reducible cubic, it is given by the pencil of lines based at 1, this follows from Bezout's theorem. The cyclic ordering of 2, $\dots, 6$ given by the lines of the successive reducible cubics is the same as the cyclic ordering of these points on the conic that they determine. The mutual cyclic orderings of 2, $\dots, 6$ given on one hand by the conic 25364, on the other hand by the pencil of lines based at 1, are displayed in the marked diagrams of Figure 12, where the circles represent the conics. Let us now consider together all six pencils of cubics with nodes at 1, $\dots, 6$. If we take 1 in one of the zones E, F, G exterior to the conic 25364, then we note that 2 lies inside of the conic determined by the other five points. So, up to the action of $\mathcal{S}_6$, we may assume that 1 is interior to a conic 25364. There are four lists of six pencils,




Figure 1: The 36 zones

denoted by $\alpha, \beta, \gamma, \delta$, see Figure 13. For a given list, each choice of a point $1, \dots, 6$ gives rise to a diagram, it turns out that all six unmarked diagrams of a list are identical, and each list corresponds to a different diagram, see Figure 14. $\square$

Rational cubics and pencils of rational cubics were used in [1]-[3] to study the topology of real algebraic M -curves of degree 9. The combinatoric of generic pencils of cubics was studied in [4], for the particular case where eight of the base points lie in convex position.




Figure 2: cremona transformations for zones A and B




Figure 3: cremona transformations for zones C and D




Figure 4: cremona transformations for zones E and F




Figure 5: cremona transformations for zone G

$1 \in A, cr(145)$	$12 \cup 36$	13462	$13 \cup 26$	$16 \cup 23$	16532
$1 \in B, cr(145)$	$12 \cup 36$	31426	$13 \cup 26$	$16 \cup 32$	25163
$1 \in C, cr(136)$	$12 \cup 45$	$15 \cup 42$	41562	41523	$14 \cup 25$
$1 \in D, cr(145)$	$12 \cup 36$	31246	$13 \cup 26$	$16 \cup 32$	16352
$1 \in E, cr(136)$	$12 \cup 45$	$15 \cup 24$	41562	41523	$14 \cup 25$
$1 \in F, cr(145)$	$12 \cup 36$	34126	$13 \cup 26$	$16 \cup 23$	32156
$1 \in G, cr(136)$	$12 \cup 45$	$15 \cup 24$	41562	41532	$14 \cup 25$

Figure 6: The seven pencils of conics

$1 \in A$	$12 \cup 51436$	$15 \cup 31264$	$13 \cup 51624$	$16 \cup 31452$	$14 \cup 21653$
$1 \in B$	$12 \cup 31645$	$15 \cup 31642$	$13 \cup 51642$	$16 \cup 31425$	$14 \cup 31625$
$1 \in C$	$12 \cup 15364$	$15 \cup 31264$	$13 \cup 51264$	$16 \cup 21453$	$14 \cup 21653$
$1 \in D$	$12 \cup 31654$	$15 \cup 31624$	$13 \cup 51642$	$16 \cup 31425$	$14 \cup 31652$
$1 \in E$	$12 \cup 15364$	$15 \cup 12346$	$13 \cup 12546$	$16 \cup 12453$	$14 \cup 12653$
$1 \in F$	$12 \cup 16453$	$15 \cup 31642$	$13 \cup 16425$	$16 \cup 13524$	$14 \cup 13526$
$1 \in G$	$12 \cup 14635$	$15 \cup 12436$	$13 \cup 12456$	$16 \cup 12453$	$14 \cup 12635$

Figure 7: The seven pencils of cubics




Figure 8: pencils with node in 1 for zones A and B




Figure 9: pencils with node in 1 for zones C and D




Figure 10: pencils with node in 1 for zones E and F




Figure 11: pencils with node in 1 for zone G




Figure 12: Marked diagrams for $1 \in A, \dots, G$, the circle represents the conic

$\alpha : 1 \in A$

$1 < 25364$	$12 \cup 51436$	$15 \cup 31264$	$13 \cup 51624$	$16 \cup 31452$	$14 \cup 21653$
$2 < 51436$	$25 \cup 31264$	$21 \cup 25364$	$24 \cup 21653$	$23 \cup 51624$	$26 \cup 31452$
$3 < 51624$	$35 \cup 31264$	$31 \cup 25364$	$36 \cup 31452$	$32 \cup 51436$	$34 \cup 21653$
$4 < 21653$	$42 \cup 51436$	$41 \cup 25364$	$46 \cup 31452$	$45 \cup 31264$	$43 \cup 51624$
$5 < 31264$	$53 \cup 51624$	$51 \cup 25364$	$52 \cup 51436$	$56 \cup 31452$	$54 \cup 21653$
$6 < 31452$	$63 \cup 51624$	$61 \cup 25364$	$64 \cup 21653$	$65 \cup 31264$	$62 \cup 51436$

$\beta : 1 \in B$

$1 < 25364$	$12 \cup 31645$	$15 \cup 31642$	$13 \cup 51642$	$16 \cup 31425$	$14 \cup 31625$
$2 > 31645$	$23 \cup 51642$	$21 \cup 25364$	$26 \cup 31425$	$24 \cup 31625$	$25 \cup 31642$
$3 > 51642$	$35 \cup 31642$	$31 \cup 25364$	$36 \cup 31425$	$34 \cup 31625$	$32 \cup 31645$
$4 < 31625$	$43 \cup 51642$	$41 \cup 25364$	$46 \cup 31425$	$42 \cup 31645$	$45 \cup 31642$
$5 < 31642$	$53 \cup 51642$	$51 \cup 25364$	$56 \cup 31425$	$54 \cup 31625$	$52 \cup 31645$
$6 > 31425$	$63 \cup 51642$	$61 \cup 25364$	$64 \cup 31625$	$62 \cup 31645$	$65 \cup 31642$

$\gamma : 1 \in C$

$1 < 25364$	$12 \cup 15364$	$15 \cup 31264$	$13 \cup 51264$	$16 \cup 21453$	$14 \cup 21653$
$2 > 15364$	$21 \cup 25364$	$25 \cup 31264$	$23 \cup 51264$	$26 \cup 21453$	$24 \cup 21653$
$3 > 51264$	$35 \cup 31264$	$31 \cup 25364$	$32 \cup 15364$	$36 \cup 21453$	$34 \cup 21653$
$4 < 21653$	$42 \cup 15364$	$41 \cup 25364$	$46 \cup 21453$	$45 \cup 31264$	$43 \cup 51264$
$5 < 31264$	$53 \cup 51264$	$51 \cup 25364$	$52 \cup 15364$	$56 \cup 21453$	$54 \cup 21653$
$6 > 21453$	$62 \cup 15364$	$61 \cup 25364$	$64 \cup 21653$	$65 \cup 31264$	$63 \cup 51264$

$\delta : 1 \in D$

$1 < 25364$	$12 \cup 31654$	$15 \cup 31624$	$13 \cup 51642$	$16 \cup 31425$	$14 \cup 31652$
$2 < 31654$	$23 \cup 51642$	$21 \cup 25364$	$26 \cup 31425$	$25 \cup 31624$	$24 \cup 31652$
$3 > 51642$	$35 \cup 31624$	$31 \cup 25364$	$36 \cup 31425$	$34 \cup 31652$	$32 \cup 31654$
$4 > 31652$	$43 \cup 51642$	$41 \cup 25364$	$46 \cup 31425$	$45 \cup 31624$	$42 \cup 31654$
$5 > 31624$	$53 \cup 51642$	$51 \cup 25364$	$56 \cup 31425$	$52 \cup 31654$	$54 \cup 31652$
$6 > 31425$	$63 \cup 51642$	$61 \cup 25364$	$64 \cup 31652$	$62 \cup 31654$	$65 \cup 31624$

Figure 13: The four lists of six pencils




Figure 14: The mutual cyclic orderings of the points on the conics and in the pencils of lines

References

- [1] S. Fiedler-Le Touzé: *Cubics as tool to study the topology of M-curves of degree 9 in $\mathbb{R}P^2$* , Journal London Math. Soc.(2) 66 (2002) 86-100.
- [2] S. Fiedler-Le Touzé: *M-curves of degree 9 with deep nests*, Journal London Math. Soc. (2009) 79(3), 649-662
- [3] S. Fiedler-Le Touzé: *M-curves of degree 9 with three nests*, arXiv [math.AG] 0806.4446v2, Sept. 2010
- [4] S. Fiedler-Le Touzé: *Pencils of cubics with eight base points lying in convex position in $\mathbb{R}P^2$* , arXiv [math AG] 1012.2679v2, last update in Sept. 2012.

severine.fiedler@live.fr